arXiv:1512.05780v1 [hep-th] 17 Dec 2015

Higher Dimensional Curved Domain Walls on Kähler Surfaces

Fiki T. Akbar[♯], Bobby E. Gunara[♯], Rio N. Wijaya[♯],
Flinn C. Radjabaycolle[♯,♭]

[♯] *Theoretical Physics Laboratory*
Theoretical High Energy Physics and Instrumentation Research Group,
Faculty of Mathematics and Natural Sciences,
Institut Teknologi Bandung
Jl. Ganesha no. 10 Bandung, Indonesia, 40132
and
[♭] *Departement of Physics*
Faculty of Mathematics and Natural Sciences,
Cendrawasih University
Jl. Kampwolker Kampus Uncen Baru Waena-Jayapura 99351, Indonesia
email: ftakbar@fi.itb.ac.id, bobby@fi.itb.ac.id

Abstract

In this paper we study some aspects of curved BPS-like domain walls in higher dimensional gravity theory coupled to scalars where the scalars span a complex Kähler surface with scalar potential turned on. Assuming that a fake superpotential has a special form which depends on Kähler potential and a holomorphic function, we prove that BPS-like equations have a local unique solution. Then, we analyze the vacuum structure of the theory including their stability using dynamical system and their existence in ultraviolet-infrared regions using renormalization group flow.

1 Introduction and Main Results

Topological defects such as domain walls in gravity-scalar systems have been studied over the last twenty years since they might provide some tools to explore the correspondence between gravitational theory in anti-de Sitter (AdS) spacetime and a conformal field theory (CFT) in flat Minkowskian spacetime (referred to as AdS/CFT correspondence) both in the context of the context of supersymmetric theories and in the the context of non-supersymmetric theories. In the first situation, there has been a lot of study investigating these walls in five dimensional supergravity, see for example [1, 2, 3, 4], while in the latter, some studies have been focused on higher dimensional models, see for example, [5, 6].

Apart from the above application, our interest here is to investigate the nature of curved domain walls of a higher dimensional gravity-scalar system with scalar potential turned on where the scalars are complex and span a two dimensional Kähler surface. These walls are described by an ansatz metric on which the codimension-one sliced spacetime has a cosmological constant, say Λ_D. In particular, we consider a model in which the scalar potential can be expressed in terms of a fake superpotential $\mathcal{W}(z, \bar{z})$ and its derivative (see eq. (2.15)) such that the dynamics of the walls are governed by first order differential equations called Bogomol'nyi-Prasad-Sommerfeld (BPS)-like equations (or gradient flow equations).

Our next step is to show that these BPS-like equations have a unique local solution. This can be achieved by rewriting BPS-like equations into the form of beta functions describing a renormalization group (RG) flow in the CFT picture. By taking a special case where $\mathcal{W}(z, \bar{z})$ depends on a real C^{k+2} Kähler potential $K(z, \bar{z})$ for integer $k \geq 1$ and a holomorphic function $W(z)$ (see eq. (3.4)), we can then construct a local C^k Lipshitz function with respect to the scalars $(z, \bar{z})$ and further show that there exists a unique local solution of BPS-like equations.

Second, our analysis on critical points of the fake superpotential $\mathcal{W}(z, \bar{z})$ shows that there are possibly three types of them, namely local minima, saddle points, and degenerate points. Each of them are related to vacuums (ground states) of the theory which can be viewed as critical points of the scalar potential. The ambient spacetimes at the vacuums become Ricci-flat or Einstein with negative cosmological constant. In the first case, the sliced spacetime should also Ricci-flat ($\Lambda_D = 0$) in order to evade ambient spacetime singularity, while in the latter case, Λ_D could be arbitrary.

If the model is "fake supersymmetric" introduced in [5, 7, 8] with $\Lambda_D = 0$, then gravitational stability further requires that the sliced spacetime should be flat Minkowskian and the ambient spacetimes simplify to maximally symmetric spacetimes, namely flat Minkowskian or maximally symmetric AdS spacetime. This conclusion is true for general form of the metric of Kähler surfaces and arbitrariness of the fake superpotential $\mathcal{W}(\phi)$. But, it becomes unworkable for $\Lambda_D \neq 0$ in general because so far, the gravitational stability analysis using Witten-Nester energy functional, for example, in [8] does not cover this case.

Finally, we conclude our analysis on gradient flow equations and RG flows at the ground states as follows. In Ricci-flat domain walls ($\Lambda_D = 0$), the vacuums can be defined in the asymptotic region and can be categorized into ultraviolet (UV) vacuums in high energy level and infrared (IR) vacuums in low energy level. In the case of Einstein domain walls, such a setup, however, could not always be well-defined in general. A way to overcome this problem is to modify the assumption that this theory can only be defined on a non-compact interval $I \subseteq \mathbb{R}$ such that the IR regions can be defined in the limit of $\mathrm{Inf}(I)$, while the UV regions are in the limit of $\mathrm{Sup}(I)$. In addition, we get the same conclusions for both cases, namely in the UV regions, all types of ground states are allowed, whereas in the IR regions, only stable local minima are permitted.

The organization of this paper can be mentioned as follows. In Section 2 we give a short discussion about domain walls of a higher dimensional gravity-scalar system where the scalars can be viewed as the complex coordinates of a Kähler surface. We show the local existence and the uniqueness of BPS-like equations in Section 3. The discussion of vacuum structure of the theory is in Section 4. We discuss the stability of vacuums in the context of dynamical system using BPS-like equations and their classification using

beta functions in Section 5. In the end, we apply the general analysis to a simple model in which the Kähler metric is flat and the holomorphic function $W(z)$ has the quadratic form.

2 $(D+1)$-Dimensional Domain Walls on Kähler Surfaces

In this section we shortly discuss a higher dimensional gravity theory coupled with a complex scalar where the scalar spans a Kähler surface explained below. Our setting here is inspired by our previous work on flat and curved domain walls of $N = 1$ chiral supergravity in four dimensions [9, 10].

Let $\left(\mathcal{M}^{D+1}, g_{\mu\nu}\right)$ be a spacetime manifold in $D+1$ dimensions equipped with metric tensor $g_{\mu\nu}$ where $\mu, \nu = 0, 1, \ldots, D$. In this paper, we are using the signature $(-, +, \ldots, +)$. The action of the Einstein-Complex Scalar system in $D+1$ dimensions has the form

$$S = \int d^{D+1}x \sqrt{-g} \left(\frac{1}{2\kappa^2} R - g_{z\bar{z}} \partial_\mu z \partial^\mu \bar{z} - V(z, \bar{z}) \right) , \tag{2.1}$$

where the above quantities can be mentioned as follows. The quantity $\kappa \equiv 1/M_P$ with M_P is Planck mass, $g \equiv \det\left(g_{\mu\nu}\right)$, and R is a scalar Ricci of $\mathcal{M}^{D+1}$. The field z is a complex scalar together with its conjugate $\bar{z}$ spanned a Kähler surfacewith metric $g_{z\bar{z}} = \partial^2 K/\partial z \partial \bar{z}$ where $K(z, \bar{z})$ is Kähler potential. In particular, we assume $g_{z\bar{z}}$ to be positive definite. Finally, the function $V(z, \bar{z})$ is a real scalar potential.

Then, the field equations of motions of the theory consist of two equations, namely the Einstein field equation,

$$\frac{1}{\kappa^2} R_{\mu\nu} = g_{z\bar{z}} \left(\partial_\mu z \partial_\nu \bar{z} + \partial_\nu z \partial_\mu \bar{z} \right) + \frac{2}{D-1} g_{\mu\nu} V(z, \bar{z}) , \tag{2.2}$$

and the scalar field equations of motions,

$$\frac{g_{z\bar{z}}}{\sqrt{-g}} \partial_\mu \left(\sqrt{-g} \partial^\mu z \right) + \partial_{\bar{z}} g_{z\bar{z}} \partial_\mu z \partial^\mu \bar{z} - \partial_{\bar{z}} V(z, \bar{z}) = 0 , \tag{2.3}$$

together with their complex conjugate.

The domain walls solution can be constructed by considering the following ansatz of the spacetime metric

$$ds^2 = a^2(u) \tilde{g}_{ab}(x^a) dx^a dx^b + du^2 , \tag{2.4}$$

where $x^\mu = (x^a, u)$, $a, b = 0, 1, \ldots, D-1$ are the local coordinates defined on $\mathcal{M}$ and $a(u)$ is the warp factor. The form of sliced metric $\tilde{g}_{ab}$ determine the type of domain walls. In this paper, we consider the case of the curved domain walls on which the sliced metric $\tilde{g}_{ab}$ has to be Einstein metric in D dimensions satisfying

$$\tilde{R}_{ab} = \Lambda_D \, \tilde{g}_{ab} , \tag{2.5}$$

where Λ_D is a cosmological constant in D dimensions known as Einstein-sliced domain walls. For $\Lambda_D = 0$, the sliced geometry is Ricci-flat and in general, this is still a curved spacetime. In this case we have Ricci-flat domain walls which generalize flat domain

walls. For $\Lambda_D > 0$ case, we have de Sitter (dS)-domain walls, whereas anti-de Sitter (AdS)-domain walls require $\Lambda_D < 0$.

Next, one can compute the Ricci tensor related to the metric (2.4) which has the form

$$R_{\mu\nu} = -\tilde{g}_{ab}\delta^a_\mu\delta^b_\nu\left[(D-1)a'^2 + aa'' - \Lambda_D\right] - \delta^D_\mu\delta^D_\nu D\left(\frac{a''}{a}\right), \tag{2.6}$$

and the Ricci scalar is given by

$$R = -D\left[(D-1)\left(\frac{a'}{a}\right)^2 + 2\frac{a''}{a} - \frac{\Lambda_D}{a^2}\right], \tag{2.7}$$

where $a' \equiv da/du$.

In the rest of the paper, we consider domain wall solutions of (2.2) and (2.3) for u-dependent scalar field. Taking $z \equiv z(u)$, the equations of motions (2.2) and (2.3) can be reduced into second order ordinary differential equations with respect to u, namely

$$\frac{a''}{a} + (D-1)\left(\frac{a'}{a}\right) - \frac{\Lambda_D}{a^2} + \frac{2\kappa^2}{(D-1)}V(z,\bar{z}) = 0, \tag{2.8}$$

$$\frac{a''}{a} + \frac{2\kappa^2}{D}g_{z\bar{z}}z'\bar{z}' + \frac{2\kappa^2}{D(D-1)}V(z,\bar{z}) = 0, \tag{2.9}$$

and

$$z'' + \frac{D}{a}a'z' + g^{z\bar{z}}\partial_{\bar{z}}g_{z\bar{z}}\,z'\bar{z}' - g^{z\bar{z}}\partial_{\bar{z}}V(z,\bar{z}) = 0\,. \tag{2.10}$$

To solve (2.9) -(2.10), let us first define a set of first order flow equations

$$\frac{a'}{a} = \pm\gamma(u)\mathcal{W}(z,\bar{z})\,, \tag{2.11}$$

$$z' = \mp\frac{(D-1)}{\kappa^2\gamma(u)}g^{z\bar{z}}\partial_{\bar{z}}\mathcal{W}\,, \tag{2.12}$$

$$\bar{z}' = \mp\frac{(D-1)}{\kappa^2\gamma(u)}g^{z\bar{z}}\partial_{z}\mathcal{W}\,, \tag{2.13}$$

where $\mathcal{W} \equiv \mathcal{W}(z,\bar{z})$ is a real function (called fake superpotential) and the real valued phase factor $\gamma(u)$ is given by

$$\gamma = \sqrt{1 + \frac{\Lambda_D}{(D-1)a^2\mathcal{W}^2}}\,. \tag{2.14}$$

Equations (2.12) and (2.13) are known as gradient flow equations (or Bogomol'nyi-Prasad-Sommerfeld (BPS)-like equations). Now, we can claim that the following statement is true. Any solution of equations (2.11), (2.12) and (2.13) are also the solution of equations (2.9) and (2.10) if the following relation holds [8, 11]

$$\kappa^2 V(z,\bar{z}) = \frac{(D-1)^2}{\kappa^2\gamma^2(u)}\,g^{z\bar{z}}\partial_z\mathcal{W}\,\partial_{\bar{z}}\mathcal{W} - \frac{D(D-1)}{2}\mathcal{W}^2\,. \tag{2.15}$$

3 Local Existence and Uniqueness

In this section, we prove the local existence and the uniqueness of (2.11) and the gradient flow equations (2.12) and (2.13). Our starting point is to rewrite equations (2.11), (2.12) and (2.13) into the form

$$\begin{aligned} a\frac{dz}{da} &= -\frac{D-1}{\kappa^2\gamma^2}g^{z\bar{z}}\frac{\partial_{\bar{z}}\mathcal{W}}{\mathcal{W}}\,, \\ a\frac{d\bar{z}}{da} &= -\frac{D-1}{\kappa^2\gamma^2}g^{z\bar{z}}\frac{\partial_{z}\mathcal{W}}{\mathcal{W}}\,. \end{aligned} \tag{3.1}$$

In the context of Gravity/CFT correspondence, (3.1) can be thought of the beta functions which gives a relevant renormalization group (RG) flow of the scalar fields [1]. We will show that the local existence and the uniqueness of equations (3.1) imply the local existence and the uniqueness of equationsof original equations, (2.11), (2.12) and (2.13).

Let us write equations (3.1) into an integral form

$$\mathbf{\Phi}(a) = \mathbf{\Phi}_0 - \int_{a_0}^{a} \mathcal{F}(s, \mathbf{\Phi}(s))\, ds\,, \tag{3.2}$$

with

$$\mathbf{\Phi} = \begin{pmatrix} z \\ \bar{z} \end{pmatrix}\,, \qquad \mathcal{F}(a,\mathbf{\Phi}) = \frac{(D-1)^2}{\kappa^2}\frac{g^{z\bar{z}}a\mathcal{W}}{(D-1)a^2\mathcal{W}^2+\Lambda_D}\begin{pmatrix} \partial_{\bar{z}}\mathcal{W} \\ \partial_{z}\mathcal{W} \end{pmatrix}\,, \tag{3.3}$$

where we already used the definition of γ in equation (2.14).

For the rest of this paper, we consider a particular case where the form of $\mathcal{W}(z,\bar{z})$ is given by

$$\mathcal{W}(z,\bar{z}) \equiv e^{\kappa^2 K/2}|W(z)|\,, \tag{3.4}$$

where $W(z)$ is a holomorphic function. Assuming that the Kähler potential $K(z,\bar{z})$ is a real C^{k+2} function for integer $k \geq 1$. Then by definition, the Kähler metric is a C^k function. Since the superpotential $W(z)$ is a holomorphic function, then it is analytic, thus a C^∞ function. So, from equation (3.4), the function $\mathcal{W}$ is a function of class C^{k+2}.

Furthermore, the function $\mathcal{F}(a,\mathbf{\Phi})$ is a C^k function with respect to $\mathbf{\Phi}$. Since $k \geq 1$, then $\mathcal{F}(a,\mathbf{\Phi})$ is a locally Lipshitz function respect to second argument, hence for a real constant M, there is exist a constant L_M which depend on M such that,

$$\left\|\mathcal{F}(a,\mathbf{\Phi}) - \mathcal{F}(a,\tilde{\mathbf{\Phi}})\right\| \leq L_M \left\|\mathbf{\Phi} - \tilde{\mathbf{\Phi}}\right\|\,, \tag{3.5}$$

whenever $\|\mathbf{\Phi}\|, \|\tilde{\mathbf{\Phi}}\| \leq M$.

Define an operator

$$\mathcal{K}(\mathbf{\Phi})(a) = \mathbf{\Phi}_0 - \int_{a_0}^{a} \mathcal{F}(s, \mathbf{\Phi}(s))\, ds\,. \tag{3.6}$$

The local existence is established by showing that the operator $\mathcal{K}$ is a contraction mapping. Let $J = [a_0, a_0+\delta]$ is an interval for a small δ and U is an open set on one dimensional Kähler manifold. Also let,

$$X = \{\mathbf{\Phi} \in C(J,U) : \mathbf{\Phi}(a_0) = \mathbf{\Phi}_0,\ \|\mathbf{\Phi}\| \leq M\}\,. \tag{3.7}$$

Defining a norm on X as

$$\|\mathbf{\Phi}\|_X = \sup_{a\in J} \|\mathbf{\Phi}(a)\| \,, \tag{3.8}$$

we have then the estimate

$$\begin{aligned} \|\mathcal{K}(\mathbf{\Phi})\|_X &\leq \|\mathbf{\Phi}_0\|_X + \sup_{a\in J}\int_{a_0}^{a} \|\mathcal{F}(s,\mathbf{\Phi}(s))\|\, ds \\ &\leq \|\mathbf{\Phi}_0\|_X + \sup_{a\in J} \left(\|\mathcal{F}(0)\| + L_M\|\mathbf{\Phi}\|\right) s \\ &\leq \|\mathbf{\Phi}_0\|_X + (a_0+\delta)\left(\|\mathcal{F}(0)\| + ML_M\right) \,. \end{aligned} \tag{3.9}$$

Taking

$$\delta \leq \min\left(\frac{1}{L_M}, \frac{1}{\|\mathcal{F}(0)\| + ML_M}\right) - a_0 \,, \tag{3.10}$$

we see that $\mathcal{K}(\mathbf{\Phi})$ map X into itself. Thus, we have the following estimate

$$\begin{aligned} \|\mathcal{K}(\mathbf{\Phi}) - \mathcal{K}(\tilde{\mathbf{\Phi}})\|_X &\leq \sup_{a\in J}\int_{a_0}^{a} \|\mathcal{F}(s,\mathbf{\Phi}(s)) - \mathcal{F}(s,\tilde{\mathbf{\Phi}}(s))\|\, ds \\ &\leq \sup_{a\in J}\sup_{a_0\leq s\leq a} s\, \|\mathcal{F}(s,\mathbf{\Phi}(s)) - \mathcal{F}(s,\tilde{\mathbf{\Phi}}(s))\| \\ &\leq L_M(a_0+\delta)\|\mathbf{\Phi} - \tilde{\mathbf{\Phi}}\|_X \,. \end{aligned} \tag{3.11}$$

Since δ satisfies (3.10), then $\mathcal{K}$ is a contraction mapping on X. By contraction mapping principle, for each initial value, there exists a unique local solution of differential equation (3.1). In addition, since $\mathcal{F}(a,\mathbf{\Phi})$ is a C^k-function respect to $\mathbf{\Phi}$, it follows that the local solution of (3.1) is a C^{k+1}-function.

Using the local solution of (3.1), the projection equation (2.11) becomes an autonomous equation of a,

$$\frac{a'}{a} = \gamma(a)\mathcal{W}(\mathbf{\Phi}(a)) \,. \tag{3.12}$$

Since $\mathcal{W}$ is a C^{k+2} function and using similar way as above, then we have a unique local solution of equation (3.12). Finally, using the composition

$$z(u) = (z\circ a)(u) \,, \tag{3.13}$$

the differential equations (2.12) and (2.13) admit a unique local solution.

4 Vacuum Structure of The Theory

In this section, we investigate some aspects of critical points of scalar potential $V(z,\bar{z})$ (2.15), which is referred to as ground states or vacua of the theory, and its relation to critical points of the fake superpotential $\mathcal{W}(z,\bar{z})$. In other words, if $\mathbf{\Phi}_c \equiv (z_c,\bar{z}_c)$ is a critical point of $\mathcal{W}(z,\bar{z})$, then

$$\partial_z\mathcal{W}(\mathbf{\Phi}_c) = \frac{\mathcal{W}(\mathbf{\Phi}_c)}{2W(z_c)}\nabla_z W(\mathbf{\Phi}_c) = 0 \,, \tag{4.1}$$

where $\nabla_z W(\mathbf{\Phi}_c) \equiv \frac{dW}{dz}(z_c) + \kappa^2 K_z(\mathbf{\Phi}_c)W(z_c)$.

4.1 Critical Points of $\mathcal{W}(z,\bar{z})$

For the case at hand, namely (3.4), we can write down the eigenvalues and the determinant of Hessian matrix of $\mathcal{W}(z,\bar{z})$ evaluated at $\mathbf{\Phi}_c$ as

$$\begin{aligned} \lambda_{\pm}^{\mathcal{W}} &= \kappa^2 g_{z\bar{z}}(\mathbf{\Phi}_c)\mathcal{W}(\mathbf{\Phi}_c) \pm 2|\partial_z^2\mathcal{W}(\mathbf{\Phi}_c)|\,, \qquad (4.2)\\ \det H_{\mathcal{W}} &= \kappa^4 g_{z\bar{z}}^2(\mathbf{\Phi}_c)\mathcal{W}^2(\mathbf{\Phi}_c) - 4|\partial_z^2\mathcal{W}(\mathbf{\Phi}_c)|^2\,, \qquad (4.3)\end{aligned}$$

where

$$\partial_z^2\mathcal{W}(\mathbf{\Phi}_c) = \frac{\kappa^2\mathcal{W}(\mathbf{\Phi}_c)}{2W(z_c)}\left[K_{zz}(\mathbf{\Phi}_c)W(\mathbf{\Phi}_c) + K_z(\mathbf{\Phi}_c)\frac{dW}{dz}(z_c) + \frac{1}{\kappa^2}\frac{d^2W}{dz^2}(z_c)\right]\,. \qquad (4.4)$$

The point $\mathbf{\Phi}_c$ is said to be a local minimum if all eigenvalues in (4.2) are positive, hence

$$\frac{\kappa^2}{2}g_{z\bar{z}}(\mathbf{\Phi}_c)\mathcal{W}(\mathbf{\Phi}_c) > |\partial_z^2\mathcal{W}(\mathbf{\Phi}_c)|\,, \qquad (4.5)$$

while a saddle point requires

$$\frac{\kappa^2}{2}g_{z\bar{z}}(\mathbf{\Phi}_c)\mathcal{W}(\mathbf{\Phi}_c) < |\partial_z^2\mathcal{W}(\mathbf{\Phi}_c)|\,, \qquad (4.6)$$

Moreover, $\mathbf{\Phi}_c$ is a degenerate critical point if

$$\frac{\kappa^2}{2}g_{z\bar{z}}(\mathbf{\Phi}_c)\mathcal{W}(\mathbf{\Phi}_c) = |\partial_z^2\mathcal{W}(\mathbf{\Phi}_c)|\,. \qquad (4.7)$$

In general, there is no local maximum point in the model.

4.2 Vacuums

Now, we can discuss the vacuum structure of the theory which can be viewed as the critical points of the scalar potential $V(z,\bar{z})$. The results here generalize our previous results of flat and curved BPS domain walls in four dimensions [9, 10].

The eigenvalues and the determinant of the Hessian matrix of $V(z,\bar{z})$ evaluated at $\mathbf{\Phi}_c$ can be written down as

$$\begin{aligned} \lambda_{\pm}^{V} &= \frac{2}{\kappa^4\gamma^2}(D-1)^2 g^{z\bar{z}}\left(\mathbf{\Phi}_c\right)\left|\partial_z^2\mathcal{W}\left(\mathbf{\Phi}_c\right)\right|^2 + (D-1)\left[\frac{D-1}{2\gamma^2} - D\right] g_{z\bar{z}}\left(\mathbf{\Phi}_c\right)\mathcal{W}^2\left(\mathbf{\Phi}_c\right)\\ &\quad \pm\frac{2}{\kappa^2}(D-1)\left[D - \frac{D-1}{\gamma^2}\right]\mathcal{W}\left(\mathbf{\Phi}_c\right)\left|\partial_z^2\mathcal{W}\left(\mathbf{\Phi}_c\right)\right|\,, \qquad (4.8)\\ \det H_V &= \frac{4\left(g^{z\bar{z}}\left(\mathbf{\Phi}_c\right)\right)^2}{\kappa^8\gamma^4}(D-1)^4\left|\partial_z^2\mathcal{W}\left(\mathbf{\Phi}_c\right)\right|^4 + \frac{1}{4}(D-1)^2\left[\frac{D-1}{2\gamma^2} - D\right]^2 g_{z\bar{z}}^2\left(\mathbf{\Phi}_c\right)\mathcal{W}^4\left(\mathbf{\Phi}_c\right)\\ &\quad -\frac{2}{\kappa^4}(D-1)^2\left[\left(\frac{D-1}{\gamma^2} - D\right)^2 + D^2\right]\mathcal{W}^2\left(\mathbf{\Phi}_c\right)\left|\partial_z^2\mathcal{W}\left(\mathbf{\Phi}_c\right)\right|^2\,. \qquad (4.9)\end{aligned}$$

Our analysis on (4.8) and (4.9) concludes that a local minimum of the scalar potential (or a stable vacuum) exists if

$$\left|\partial_z^2\mathcal{W}\left(\mathbf{\Phi}_c\right)\right| > \frac{\kappa^2 g_{z\bar{z}}\left(\mathbf{\Phi}_c\right)\mathcal{W}\left(\mathbf{\Phi}_c\right)}{2(D-1)}\left[D\left(2\gamma^2-1\right)+1\right]\,, \qquad (4.10)$$

whereas a local maximum (or an unstable vacuum) requires

$$|\partial_z^2 \mathcal{W}(\mathbf{\Phi}_c)| < \frac{\kappa^2}{2} g_{z\bar{z}}(\mathbf{\Phi}_c)\, \mathcal{W}(\mathbf{\Phi}_c) \ . \tag{4.11}$$

The existence of a saddle point (or a saddle vacuum) demands

$$\frac{\kappa^2}{2} g_{z\bar{z}}(\mathbf{\Phi}_c)\, \mathcal{W}(\mathbf{\Phi}_c) < |\partial_z^2 \mathcal{W}(\mathbf{\Phi}_c)| < \frac{\kappa^2 g_{z\bar{z}}(\mathbf{\Phi}_c)\, \mathcal{W}(\mathbf{\Phi}_c)}{2(D-1)} \left[D\left(2\gamma^2 - 1\right) + 1\right] \ , \tag{4.12}$$

while a critical point is said to be degenerate if

$$|\partial_z^2 \mathcal{W}(\mathbf{\Phi}_c)| = \frac{\kappa^2 g_{z\bar{z}}(\mathbf{\Phi}_c)\, \mathcal{W}(\mathbf{\Phi}_c)}{2(D-1)} \left[D\left(2\gamma^2 - 1\right) + 1\right] \ , \tag{4.13}$$

$$|\partial_z^2 \mathcal{W}(\mathbf{\Phi}_c)| = \frac{\kappa^2}{2} g_{z\bar{z}}(\mathbf{\Phi}_c)\, \mathcal{W}(\mathbf{\Phi}_c) \ . \tag{4.14}$$

From equations (4.5), (4.6), (4.11) and (4.12), any local minima of $\mathcal{W}(z,\bar{z})$ are mapped into local maxima of the scalar potential $V(z,\bar{z})$ and saddle point of $\mathcal{W}(z,\bar{z})$ are mapped into saddles or local minima of the scalar potential $V(z,\bar{z})$. Moreover the equation (4.14) comes naturally from the fact that degenerate critical points of $\mathcal{W}$ are mapped into degenerate points of $V(z,\bar{z})$. Since both Hessian matrix $H_{\mathcal{W}}$ and H_V evaluated at the points vanish, these special points are referred to as *intrinsic* degenerate critical points of $V(z,\bar{z})$.

4.3 Ambient Spacetimes At Vacuums

Since the gradient of $\mathcal{W}(z,\bar{z})$ vanishes at $\mathbf{\Phi}_c$, the ambient spacetime $\mathcal{M}^{D+1}$ becomes Einstein with negative scalar curvature whose Ricci tensor is given by

$$R_{\mu\nu} = -D\ \mathcal{W}(\mathbf{\Phi}_c)\, g_{\mu\nu} \ , \tag{4.15}$$

while the warped factor $a(u)$ has the form

$$a(u) = \frac{1}{2}\left[\left(a_0 + \sqrt{a_0^2 + \frac{\Lambda_D}{(D-1)\mathcal{W}^2(\mathbf{\Phi}_c)}}\right) e^{\mathcal{W}(\mathbf{\Phi}_c)u} - \frac{\Lambda_D}{(D-1)\mathcal{W}(\mathbf{\Phi}_c)^2} \frac{e^{-\mathcal{W}(\mathbf{\Phi}_c)u}}{\left(a_0 + \sqrt{a_0^2 + \frac{\Lambda_D}{(D-1)\mathcal{W}^2(\mathbf{\Phi}_c)}}\right)}\right] \ , \tag{4.16}$$

with a_0 is a non-zero real constant. So, there are two possibilities of the ambient spacetimes at vacuums, namely the Ricci-flat spacetimes for $\mathcal{W}(\mathbf{\Phi}_c) = 0$ and AdS_{D+1} spacetimes for $\mathcal{W}(\mathbf{\Phi}_c) \neq 0$.

In the first case, it is necessary that the sliced geometry should also be Ricci-flat, namely $\Lambda_D = 0$, in order to evade the singularity of the warped factor $a(u)$. Ricci-flatness further implies that the vacuums are determined by the critical point of the holomorphic superpotential

$$\frac{dW}{dz}(z_c) = 0 \ . \tag{4.17}$$

since $W(z_c) = 0$. Particularly, if we consider particularly a "fake supersymmetric" model where one can introduce a set of fake supersymmetric transformations of a gravitino and a spin-$\frac{1}{2}$ fermion together with Witten-Nester energy functional [8, 7], then we have a case

as follows. In order to have "fake supersymmetric" vacuums, both ambient and sliced spacetimes become flat Minkowskian. Such a configuration is gravitationally stable since the Witten-Nester energy vanishes.

In AdS_{D+1} spacetimes, we must have non-zero $\mathcal{W}(\mathbf{\Phi}_c)$ where $\mathbf{\Phi}_c$ is the solution of the following nonholomorphic equation

$$\nabla_z W(\mathbf{\Phi}_c) \equiv \frac{dW}{dz}(z_c) + \kappa^2 K_z(\mathbf{\Phi}_c) W(z_c) = 0 \ . \tag{4.18}$$

For the case at hand, we might have several possibilities as follows. The sliced spacetime could be de Sitter ($\Lambda_D > 0$), anti-de Sitter ($\Lambda_D < 0$), and Ricci-flat ($\Lambda_D = 0$). In addition, we could have a situation where the sliced cosmological constant Λ_D is set to be $\Lambda_D = c\mathcal{W}^2(\mathbf{\Phi}_c)$ where c is a non-zero real constant. In the latter case, as $\mathcal{W}(\mathbf{\Phi}_c) \to 0$, we would regain the preceding Ricci-flat case.

It is important to mention a remark as follows. For $\Lambda_D = 0$, if the model is fake supersymmetric, then the sliced spacetime simplifies to flat Minkowskian and the ambient spacetime becomes a maximally AdS spacetime. Such a configuration is gravitationally stable. In the case of $\Lambda_D \neq 0$, it is however still unclear in general to check the gravitational stability of (2.11)-(2.13) with scalar potential (2.15) because so far, the conventional method using Witten-Nester energy functional, for example [8], does not cover this case.

5 Gradient Flows vs RG Flows

In this section, we complete our analysis on the nature of vacuums using the gradient flow equations (2.12)- (2.13) and in context of the field theory using RG flow equations (3.1) where the warped factor $a(u)$ can be viewed as an energy scale. In the case of Ricci-flat domain walls ($\Lambda_D = 0$) the vacuum structure is simpler than Einstein domain walls ($\Lambda_D \neq 0$) since we do not need an additional analysis on the warped factor $a(u)$ as we will see below.

Let us start by considering the Ricci-flat domain walls. The vacuums here are in the asymptotic region, *i.e.* $u = \pm\infty$ and can be classified into ultraviolet (UV) vacuums for $a \to +\infty$ and infrared (IR) vacuums for $a \to 0$. Expanding (2.12)- (2.13) around $\mathbf{\Phi}_c$, we then obtain the eigenvalues of the first order expansion matrix

$$\lambda^{(0)}_{\mp} = \mp\frac{(D-1)}{\kappa^2} g^{z\bar{z}}(\mathbf{\Phi}_c)\lambda^{\mathcal{W}}_{\pm} = \frac{(D-1)}{\kappa^2}\left(\mp\kappa^2\mathcal{W}(\mathbf{\Phi}_c) - 2g^{z\bar{z}}(\mathbf{\Phi}_c)|\partial_z^2\mathcal{W}(\mathbf{\Phi}_c)|\right) \ . \tag{5.1}$$

According to Lyapunov theorem, a stable flow requires that all eigenvalues of (5.1) have to be negative which follows [9]

$$|\partial_z^2\mathcal{W}(\mathbf{\Phi}_c)| > \frac{\kappa^2}{2} g_{z\bar{z}}(\mathbf{\Phi}_c)\,\mathcal{W}(\mathbf{\Phi}_c) \ , \tag{5.2}$$

flowing along local minima and the stable directions of saddles of the scalar potential (2.15). In other words, the evolution of the complex scalars $(z, \bar{z})$ is stable on the walls in the context of dynamical system. In particular, if the sliced geometry is flat Minkowskian, then we have a *complete stable* model which means that it is stable both in the context of gravitational stability and in the context of dynamical system. Other models would be *incomplete stable*.

In the case at hand, (4.16) simplifies to

$$a_{\rm rf}(u) = a_0\ e^{\mathcal{W}(\mathbf{\Phi}_c)u}\ , \tag{5.3}$$

which shows that at $u \to +\infty$, we have $a_{\rm rf} \to +\infty$ (the UV vacuums), while at $u \to -\infty$, we have $a_{\rm rf} \to 0$ (the IR vacuums). The eigenvalues of first order expansion of the beta function (3.1) are

$$\tilde{\lambda}^{(0)}_{\pm} = \frac{(D-1)}{\kappa^2}\frac{g^{z\bar{z}}(\mathbf{\Phi}_c)}{\mathcal{W}(\mathbf{\Phi}_c)}\lambda^{\mathcal{W}}_{\pm} = \frac{(D-1)}{\kappa^2}\left(\kappa^2 \pm 2\frac{g^{z\bar{z}}(\mathbf{\Phi}_c)}{\mathcal{W}(\mathbf{\Phi}_c)}|\partial_z^2\mathcal{W}(\mathbf{\Phi}_c)|\right)\ . \tag{5.4}$$

In the UV region, we require at least one of the eigenvalues (5.4) to be positive. So, the RG flow is stable along $\tilde{\lambda}^{(0)}_{+}$, but it might be unstable in the direction of $\tilde{\lambda}^{(0)}_{-}$ if $\tilde{\lambda}^{(0)}_{-} \leq 0$ which implies that the flow fails to depart the UV region. In conclusion, we have here all type of vacuums, namely local maxima (unstable), local minima (stable), saddles (stable), and bifurcation points (undetermined). On the other hand, in the IR region the RG flow is stable along the direction of the negative eigenvalue of $\tilde{\lambda}^{(0)}_{-}$, but it might be failed to approach vacuums if $\tilde{\lambda}^{(0)}_{-} \geq 0$. Thus, there are no local maxima and bifurcation points.

In Einstein domain walls, the eigenvalues of the first order expansion of gradient flow equations (2.12)- (2.13) are

$$\lambda^{(\Lambda_D)}_{\mp} = \mp\frac{(D-1)}{\kappa^2\gamma^2}g^{z\bar{z}}(\mathbf{\Phi}_c)\lambda^{\mathcal{W}}_{\pm} = \frac{(D-1)}{\kappa^2\gamma^2}\left(\mp\kappa^2\mathcal{W}(\mathbf{\Phi}_c) - 2g^{z\bar{z}}(\mathbf{\Phi}_c)|\partial_z^2\mathcal{W}(\mathbf{\Phi}_c)|\right)\ , \tag{5.5}$$

whereas the eigenvalues of first order expansion of the beta function (3.1) have the form

$$\tilde{\lambda}^{(\Lambda_D)}_{\pm} = \frac{(D-1)}{\kappa^2\gamma^2}\frac{g^{z\bar{z}}(\mathbf{\Phi}_c)}{\mathcal{W}(\mathbf{\Phi}_c)}\lambda^{\mathcal{W}}_{\pm} = \frac{(D-1)}{\kappa^2\gamma^2}\left(\kappa^2 \pm 2\frac{g^{z\bar{z}}(\mathbf{\Phi}_c)}{\mathcal{W}(\mathbf{\Phi}_c)}|\partial_z^2\mathcal{W}(\mathbf{\Phi}_c)|\right)\ . \tag{5.6}$$

Comparing (5.5) - (5.6) with (5.1) and (5.4), it is easily to see that we will get the same results as the previous Ricci-flat case. These results, however, need the analysis on the warped factor (4.16) in order to have a well-defined field theory on the boundaries. First, if we want the UV vacuums ($a \to +\infty$) and the IR vacuums (($a \to 0$)) in the limit of $u \to \pm\infty$, then the sliced spacetime should be a weak Ricci-flat with small cosmological constant, namely $\Lambda_D \approx 0$, or we have a case of $\Lambda_D \ll \mathcal{W}^2(\mathbf{\Phi}_c)$. Second, we modify the theory such that a well-defined field theory could only be defined on the interval $-|\Lambda_D|^{-1} \leq u \leq |\Lambda_D|^{-1}$. The UV vacuums can be introduced in the limit of $u \to |\Lambda_D|^{-1}$, while the IR vacuums are in the limit of $u \to -|\Lambda_D|^{-1}$ where $a(|\Lambda_D|^{-1}) > a(-|\Lambda_D|^{-1})$. Note that our analysis fails for the case of $|\Lambda_D| \to +\infty$.

6 A Simple Model

In this section we apply the previous general setup to a simple model where the Kähler space is flat with potential $K(z,\bar{z}) = z\bar{z}$ and the superpotential has the quadratic form, namely $W(z) = \alpha_0 + \alpha_1 z + \alpha_2 z^2$ where α_0, α_1, α_2 are real constants. Here, we assume the origin to be a critical point of this model which implies that $K_z(0) = 0$. Then, (4.2)

and (4.8) simplify to

$$\lambda^{\mathcal{W}}_{\pm} = \kappa^2|\alpha_0| \pm 2|\alpha_2| \ , \tag{6.1}$$

$$\lambda^{V}_{\pm} = \frac{2}{\kappa^4\gamma^2}(D-1)^2|\alpha_2|^2 + (D-1)\left[\frac{D-1}{2\gamma^2} - D\right]|\alpha_0|^2 \pm \frac{2}{\kappa^2}(D-1)\left[D - \frac{D-1}{\gamma^2}\right]|\alpha_0||\alpha_2| \ , \tag{6.2}$$

$$\tag{6.3}$$

and the scalar curvature of the ambient spacetime equals $-D(D+1)|\alpha_0|$.

For $\alpha_0 = 0$, we have a complete stable Minkowskian ground state with $\Lambda_D = 0$, while an AdS ground state requires $\alpha_0 \neq 0$ for any Λ_D. In the latter case, we have an AdS vacuum for $\alpha_2 = 0$ which is unstable in the context dynamical system, but for $\Lambda_D = 0$ it is easy to check that this vacuum is gravitationally stable. In other words, this vacuum is incomplete stable. The anaysis on RG flow shows that this vacuum exists only in the UV region.

Acknowledgements

Our research is supported by Riset KK ITB 2015 and Riset Desentralisasi DIKTI-ITB 2015.

References

[1] D.Z. Freedman, S.S. Gubser, K. Pilch and N.P. Warner, "Renormalization group flows from holography-supersymmetry and a c-theorem", *Adv. Theor. Math. Phys.* **3** (1999) 363, hep-th/9904017.

[2] A. Ceresole, G. Dall'Agata, R. Kallosh, and A. van Proeyen, *Hypermultiplets, Domain Walls and Supersymmetric Attractor*, Phys. Rev. **D64** (2001) 104006, hep-th/0104056.

[3] G. L. Cardoso, G. Dall'Agata, and D. Lüst, "Curved BPS domain walls and RG flow in five dimensions", *J. High Energy Phys.* **0203** (2002) 044, hep-th/0201270.

[4] G. L. Cardoso and D. Lüst, "The holographic RG flow in a field theory on a curved background", *J. High Energy Phys.* **0209** (2002) 028, hep-th/0207024.

[5] K. Skenderis and P.K. Townsend, "Gravitational Stability and Renormalization Group Flow", *Phys. Lett.* **B468** (1999) 46, hep-th/9909070.

[6] R. C. Meyrs and A. Singh, "Comments on Holographic Entanglement Entropy and RG Flows", *J. High Energy Phys.* **1204** (2012) 122, hep-th/1202.2068.

[7] P. K. Townsend, "POSITIVE ENERGY AND THE SCALAR POTENTIAL IN HIGHER DIMENSIONAL (SUPER) GRAVITY THEORIES", *Phys. Lett.* **B148** (1984), 55.

[8] D. Z. Freedman, C. Núñez, M. Schnabl, and K. Skenderis, "Fake supergravity and domain wall stability", *Phys. Rev.* **D69** (2004), 104027.

[9] B. E. Gunara, F.P. Zen, and Arianto, "BPS Domain Walls and Vacuum Structure of N=1 Supergravity Coupled to A Chiral Multiplet", *J. Math. Phys.* **48** (2007), 053505, hep-th/0612017.

[10] B. E. Gunara and F.P. Zen, "Deformation of Curved BPS Domain Walls and Supersymmetric Flows on 2d Khler-Ricci Solito", *Commun. Math. Phys.* **287** (2009), 849, hep-th/0801.0896.

[11] O. DeWolfe, D. Z. Freedman, S. S. Gubser, and A. Karch, "Modelling the fifth dimension with scalars and gravity", *Phys. Rev.* **D62** (2000), 046008.